

## 2015 Western Regional Meeting 224

### Nickel-catalyzed activation of amides and simple esters

**Liana Hie**<sup>2</sup>, [lhie@chem.ucla.edu](mailto:lhie@chem.ucla.edu), Neil K. Garg<sup>1</sup>, Noah F. Fine Nathel<sup>4</sup>, Tejas Shah<sup>5</sup>, Emma Baker<sup>2</sup>, Xin Hong<sup>6</sup>, Yun-Fang Yang<sup>2</sup>, Peng Liu<sup>3</sup>, Kendall N. Houk<sup>4</sup>. (1) UCLA Chemistry, Los Angeles, California, United States (2) Box 951569, UCLA Dept of Chem BioChem, Los Angeles, California, United States (3) Department of Chemistry, University of Pittsburgh, Los Angeles, California, United States (4) UCLA, Los Angeles, California, United States


This presentation will focus on our recent efforts to develop reactions of acyl electrophiles using nickel catalysis. The methodologies proceed under mild reaction conditions and allow for the interconversion of amides and esters. These studies are expected to fuel the further use of non-precious metal catalysis for the construction of C–heteroatom and C–C bonds.

## 2015 Western Regional Meeting 225

### Nickel-catalyzed asymmetric reductive cross-coupling between heteroaryl iodides and $\alpha$ -chloronitriles

**Nathaniel T. Kadunce**, [nkadunce@caltech.edu](mailto:nkadunce@caltech.edu). Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California, United States

Ni-catalyzed asymmetric reductive cross-coupling provides an attractive and powerful means to access tertiary stereocenters. However these methods had previously been limited to benzylic alkyl partners and were not amenable to heterocyclic substrates. To expand the utility of this class of reactions, the Ni-catalyzed asymmetric cross-electrophile coupling of heteroaryl iodides and  $\alpha$ -chloronitriles has been developed. This method furnishes enantioenriched  $\alpha,\alpha$ -disubstituted nitriles from simple organohalide building blocks. The reaction tolerates a variety of heterocyclic coupling partners, including pyridines, pyrimidines, quinolines, thiophenes, and piperidines under mild conditions enabled by a novel ligand scaffold. The products can be derivatized to a range of synthetically useful functionality.



## 2015 Western Regional Meeting 226

### Review of biofuels and biofuels-related technology patents and patent applications